

## Ectopic kidney urothelial carcinoma with lower rectum adenocarcinoma: A case report

Hamdan S Alaamri<sup>1</sup>, Fawaz Al Thobaiti<sup>1</sup>, Mohammad Al Adwani<sup>1</sup>, Hama A Abdelaziz<sup>1</sup>, Amjad Ahmed Althagafi<sup>1</sup>, Faisal S Alrubaei<sup>2</sup>, Sara A Assiri<sup>3</sup>, Raad M M Althaqafi<sup>2\*</sup>

### To Cite:

Alaamri HS, Al-Thobaiti F, Al-Adwani M, Abdelaziz HA, Althagafi AA, Alrubaei FS, Assiri SA, Althaqafi RMM. Ectopic kidney urothelial carcinoma with lower rectum adenocarcinoma: A case report. Medical Science, 2022, 26, ms87e2112.  
doi: <https://doi.org/10.54905/disssi/v26i121/ms87e2112>

### Authors' Affiliation:

<sup>1</sup>Surgery department, Alhada Military Hospital, Taif, Saudi Arabia  
<sup>2</sup>College of Medicine, Taif University, Taif, Saudi Arabia  
<sup>3</sup>Otolaryngology-Head and Neck Surgery Department, King Faisal Hospital, Taif, Saudi Arabia

### \*Corresponding Author

College of Medicine, Taif University, Taif,  
Saudi Arabia  
Email: raadmatar977@gmail.com

### Peer-Review History

Received: 27 January 2022  
Reviewed & Revised: 28/January/2022 to 24/February/2022  
Accepted: 25 February 2022  
Published: 02 March 2022

### Peer-review Method

External peer-review was done through double-blind method.

URL: <https://www.discoveryjournals.org/medicalscience>



This work is licensed under a Creative Commons Attribution 4.0 International License.

### ABSTRACT

Bladder cancer (BC), one of the urothelial carcinoma (UC) and colorectal cancer (CRC) are Lynch syndrome's manifestations. In this case report, a 56 years old woman was admitted to ER with intermittent, progressive and colicky abdominal pain for 6 months associated with a change in bowel habit and bleeding per rectum, anorexia and weight loss. Rectal examination revealed a nodular, easy to bleed mass from anal verge. MRI revealed left ectopic pelvic kidney with soft tissue enhancing lesion in the lower calyx. Patient was underwent low anterior resection and left nephrectomy after which she was decided to complete on adjuvant XRT and chemotherapy. The comorbidity of UC and CRC in this patient in spite of being of low risky group being a non-smoker female has strongly prove the existence of the lynch syndrome and may replace the molecular confirmation. Therefore genetic investigation of all her relatives for presence of causative genetic abnormalities is mandatory, a project we are currently working on to enable proper medical intervention in the proper time.

**Keywords:** Ectopic kidney, UC, CRC, Woman

### 1. INTRODUCTION

Bladder cancer (BC) is one of the most frequent types of cancer encountered in the urogenital tract (Minoli et al., 2020; Vikas et al., 2020). Its responsibility for a 549,000 new cases and 200,000 death/year worldwide has makes it the 10th most common type of cancers (Bray et al., 2018). Bladder cancers are of various type including urothelial carcinoma, squamous cell carcinoma, small cell carcinoma, adenocarcinoma and sarcoma, among which carcinoma of the tract covering is the cause of most cases (Minoli et al., 2020). This urothelial carcinoma type of bladder cancer can emerge in any part of the urine passage covered with urothelium but it is confined predominantly in the lower part (bladder, urethra) and rarely in the upper tract (pelvis, ureter) (Rouprêt et al., 2015).


Smoking and male gender are among the high risk elements in addition to exposure to industrial chemicals, ethnicity and age (American Cancer Society, 2014). Of this urothelial bladder cancer, the cancer type confined to the bladder lining (non-metastatic) represents about 75% of all new cases (Kamat et al., 2016), while the metastatic type of the bladder cancer, beside other types of BC representing about 25% of cases (Robertson et al., 2017). Half of the metastatic type patients have a life expectancy less than 5 years meanwhile above 90% of non-metastatic patients life expectancy is extended after 5 years after the treatment (Knowles & Hurst 2015). Induction of a mutation in the colon germ line DNA mismatch repair (MMR) genes results in colon cancer development called the autosomal dominant hereditary cancer of the colon and rectum with minimal polyps numbers or lynch syndrome (Lynch et al., 2009), which is usually detected as either microsatellite instability in certain DNA sequences or loss of immunostaining for the respective protein (Hartmann et al., 2003).

Extra colonic cancer is a common incident with Lynch type cancer development and patients may be presented mainly with extra colonic malignancy (Roupret et al., 2008). Lynch syndrome is the genetic background of about 2–3% of all cancer cases of colon-rectum segment (Bhalla et al., 2013) and is connected with different type of neoplasms including carcinoma of the urinary passage lining, specifically of early portion the of urinary tract (Roupret et al., 2008; Mork et al., 2015). In spite of its low incidence in female, we currently reporting a case of urothelial carcinoma in the ectopic kidney's renal pelvis associated with lower rectum adenocarcinoma in a 56 years old woman.

## 2. CASE PRESENTATION


A 56 year old female patient known case of DM on medication with no past surgical history and no history of food or drug allergies was presented to ER complaining of Intermittent, progressive and colicky in nature abdominal pain for six months. With no aggravating or relieving factors, this pain was combined with a disturbance in bowel habit and bleeding per rectum, history of anorexia and weight loss of 10kg. No other systemic indications of surgical importance. On examination, patient was conscious and alert with all her vital signs being within normal. Abdomen was soft and lax, no palpable mass, no organomegaly. DRE showed a mass around 4 cm from anal verge, nodular and easy to bleed. Haematological investigations showed: WBC 6.9, HB 12.8, PLT 178, Creatinine 105, Urea 6.5. No tumor markers. Pan CT done for metastatic screening and showed no classical pulmonary or hepatic deposits. Sagittal T2W1 showed post contrast enhancement that revealed left side ectopic pelvic kidney with the lower pole soft tissue mildly enhancing mass lesion seen casting the lower calyx with heterogeneous dominantly dark signal in T2 W1 (figure 1).

Patient was sent to urology after the CT finding and they did cystoscopy for her and biopsy was taken from left renal pelvis. Patient case was discussed in the tumor committee and decision was taken for low anterior resection and left nephrectomy (figure 2). Patient underwent ultra-low anterior resection with diversion loop ileostomy with left nephroureterectomy and specimens were sent for histopathology that revealed non-invasive papillary low grade urothelial carcinoma in renal pelvis of ectopic kidney (figure 3) and papillae with fibrovascular cores lined by neoplastic cells (figure 4). This papilla showed mild architectural disorder, mitotic figure and mild nuclear pleomorphism in high power examination (figure 5). Histopathological examination of the excised rectal mass revealed; rectal invasive adenocarcinoma with neoplastic glands infiltrating the rectal wall (figure 6). Patient was discussed again postoperatively in tumor board and decision was made for adjuvant XRT and chemotherapy.




**Figure 1** (A) sagittal T2W1 and (B) axial T1W1 showing circumferential neoplastic wall thickening of mid portion of the rectum with no sizable exophytic soft tissue component (yellow arrow) denoting T2 (T2N1). (Arrow head) denoting tiny perirectal lymph nodes. Rectal carcinoma T2 N1 M0






**Figure 2** sagittal (A) T2W1 and B axial T1W1 showing post contrast enhancement that revealed left side ectopic pelvic kidney with the lower pole soft tissue mildly enhancing mass lesion seen casting the lower calyx with heterogeneous dominantly dark signal in T2 W1 ( Yellow arrow )



**Figure 3** Renal pelvis of ectopic kidney. (A) Low power showing non-invasive papillary low grade urothelial carcinoma. Papillae with fibrovascular cores lined by neoplastic cells (B) High power showing papilla with mild architectural disorder, mitotic figure and mild nuclear pleomorphism.



**Figure 5** Rectal invasive adenocarcinoma with (A) low power showing neoplastic glands infiltrating the rectal wall. (B) High power showing: characteristic intestinal type features (pencil nuclei, purple cytoplasm, rounded glands) Glands show dirty necrosis (arrow).

### 3. DISCUSSION

Colorectal carcinoma due to hereditary reasons is caused mostly by the Lynch described type of colorectal cancer being the genetic base of about 2–3% of all new colorectal carcinomas (Bhalla et al., 2013). Urothelial cancer which is the third most frequent cancer the hereditary cancer families (Lynch et al., 1990), may be encountered in some instance associated with ectopic kidney. Renal ectopic reported in this presented, case is a common congenital anomalies occurred during intra-uterine development due to failure of ascent of the kidney from the pelvis.

The worldwide incidence of renal ectopia is about 1/1000 in different ectopic locations (Gleason et al., 1994), among which pelvic renal ectopia incidence is approximately 1/2200 to 1/3000 necropsies (Oyinloye et al., 2020) therefore, the pelvic renal ectopia presented in the current case is considered a rare condition. Although renal ectopia is generally not life-threatening but in a considerable part of its occurrence, it is linked with other complications of lynch syndrome.

In the currently reported case, the concomitant occurrence of the colorectal cancer (CRC) and UC of the early portion of the urinary tract is in consistence with previous reports stating that colorectal carcinoma and UC of the early portion of the urine passage are the highest rated malignant tumor occurs in viscera that can be identified in the subset of the lynch syndrome known as Muir-Torre syndrome that was identified to have MSH2 mutation (Giardiello et al., 2014; Roupert et al., 2008; Ericson et al., 2003). The association between urothelial carcinoma and the lower CRC in this reported case may indicate the presence of MMR as a manifestation of the Lynch syndrome. Indeed, deficient mismatch repair (dMMR) reached a 15 % in sporadic CRC of Thai patients (Korphaisarn et al., 2015).

The currently investigated case is considered one of the low risk group because of being female, non-smoker and have no history of exposure to industrial chemicals (American Cancer Society, 2014). However, lynch syndrome patients commonly are not in need for the expected risk inducers to develop urothelial carcinoma (Sijmons et al., 1998; Roupert et al., 2008). Moreover, the patients of this genetic predisposition are already recognized of possess about 20% chance of growing UC in the early urinary canal in the long run (Van der et al., 2010).

Although lynch syndrome need to be confirmed by the molecular investigation, the comorbidity of UC and CRC in this patient may strongly confirm the presence of the lynch syndrome and may replace the molecular confirmation. The appearance of a single lynch syndrome's manifestation in any individual may necessitate the genetic investigation of all his relatives for the causative genetic abnormalities (loss of MMR protein or lynch syndrome), a project we are currently working on. This genetic screening of here family members may enables a proper medical intervention in the proper time.

### 4. CONCLUSION

Although the main complain of this patient was intermittent, progressive abdominal pain for 6 months with bowel habit changes and bleeding per rectum, anorexia and weight loss. Absence of the known classical tumor predisposing causes in this woman may shift the investigator thinking fare from tumor presence. However the precise clinical examination and whole body screening with CT and MRI revealed renal ectopia and presence of UC and CRC that were considered as lynch syndrome's manifestation. This concurrent occurrence of such two lynch syndrome's manifestation may be taken as substitute for the genetic screening.

#### Acknowledgments

We are extremely grateful for the Research Ethics Committee, at Al-Hada Armed Forces Hospital Taif city, Saudi Arabia for facilitating obtaining required data.

#### Informed consent

Oral and written informed consent had been got from the woman reported in this case.

#### Funding

This study has not received any external funding.

#### Conflict of interests

The authors declare that there are no conflicts of interests.

#### Data and materials availability

All data associated with this study are present in the paper.



# REFERENCES AND NOTES

1. American Cancer Society 2014. What are the Risk Factors for Bladder Cancer? <https://www.cancer.org/cancer/bladder-cancer/causes-risks-prevention/risk-factors.html>.
2. Bhalla A, Zulfiqar M, Weindel M, Vinod B Shidham. Molecular diagnostics in colorectal carcinoma. Clin Lab Med 2013; 33:835–859. PMID: 24267189 DOI: 10.1016/j.cl.2013.10.001.
3. Bray B, Ferlay J, Soerjomataram I, Siegel R L, Torre L A, Jemal A. Global Cancer Statistics. GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. CA Cancer J Clin 2018; 68:394–424. PMID: 30207593 DOI: 10.3322/caac.21492.
4. Ericson K, Halvarsson B, Nagel J, M Planck, Z Piotrowska, H Olsson, M Nilbert. Defective mismatch-repair in patients with multiple primary tumours including colorectal cancer. Eur J Cancer 2003; 39:240–248. PMID: 12509957 DOI: 10.1016/s0959-8049(02)00584-1.
5. Giardiello FM, Allen JI, Axilbund JE, C Richard Boland , Carol A Burke, Randall W Burt , James M Church , Jason A Dominitz , David A Johnson , Tonya Kaltenbach , Theodore R Levin , David A Lieberman , Douglas J Robertson , Sapna Syngal . Guidelines on genetic evaluation and management of Lynch syndrome: a consensus statement by the US Multisociety Task Force on colorectal cancer. Am J Gastroenterol 2014; 109:1159–1179. PMID: 25070057 DOI: 10.1038/ajg.2014.186.
6. Gleason PE, Kelalis PP, Husmann DA, Kramer SA. Hydronephrosis in renal ectopia: incidence, etiology and significance. J Urol 1994; 151: 1660–1 PMID: 8189592 DOI: 10.1016/s0022-5347(17)35338-7.
7. Hartmann Arndt, John C Cheville, Wolfgang Dietmaier, Ferdinand Hofstädter, Lawrence J Burgart, Hagen Blaszyk. Hereditary Nonpolyposis Colorectal Cancer Syndrome in a Patient with Urothelial Carcinoma of the Upper Urothelial Tract. Arch Pathol Lab Med 2003; 127: 60–63. PMID: 12562253 DOI: 10.5858/2003-127-e60-HNCCSI.
8. Kamat AM, Hahn NM, Efstathiou JA, Lerner SP, Malmström PU, Choi W, Guo CC, Lotan Y, Kassouf W. Bladder cancer. Lancet (London, England). 2016; 388:2796–2810. PMID: 27345655 DOI: 10.1016/S0140-6736(16)30512-8.
9. Knowles MA, Hurst CD. Molecular biology of bladder cancer: new insights into pathogenesis and clinical diversity. Nat Rev Cancer 2015; 15(1):25–41. PMID: 25533674 DOI: 10.1038/nrc3817.
10. Korphaisarn K, Pongpaibul A, Limwongse C, Roothumnon E, Klaisuban W, Nimmannit A, Jinawath A, Akewanlop C. Deficient DNA mismatch repair is associated with favorable prognosis in Thai patients with sporadic colorectal cancer. World J Gastroenterol 2015; 21(3): 926-934. PMID: 25624727 DOI: 10.3748/wjg.v21.i3.926.
11. Lynch HT, Ens JA, Lynch JF. The Lynch syndrome II and urological malignancies. J Urol 1990; 143: 24–28. PMID: 2294255 DOI: 10.1016/s0022-5347(17)39853-1.
12. Lynch HT, Lynch P M, Lanspa S J, Snyder C L, Lynch JF, Boland C R. Review of the Lynch syndrome: history, molecular genetics, screening, differential diagnosis, and medicolegal ramifications. Clin Genet 2009; 76:1–18. PMID: 19659756 DOI: 10.1111/j.1399-0004.2009.01230.x.
13. Maestro ML, Vidaurreta M, Sanz-Casla MT, Rafael S, Veganzones S, Martínez A, Aguilera C, Herranz MD, Cerdán J, Arroyo M. Role of the BRAF mutations in the microsatellite instability genetic pathway in sporadic colorectal cancer. Ann Surg Oncol 2007; 14: 1229-1236 [PMID: 17195912 DOI: 10.1245/s10434-006-9111-z.
14. Minoli M, Kiener M, Thalmann GN, Kruithof-de Julio M, Seiler R. Evaluation of Urothelial bladder Cancer in the context of molecular classification. Int J Mol Sci 2020; 21:5670. PMID: 32784716 DOI: 10.3390/ijms21165670.
15. Mork M, Hubosky SG, Roupert M. Lynch Syndrome: A Primer for Urologists and Panel Recommendations. J Urol 2015; 194:21–29. PMID: 25711197 DOI: 10.1016/j.juro.2015.02.081.
16. Oyinloye AO, Wabada S, Abubakar AM, Oyeibanji LO, Rikin CU. Wilms tumor in a left pelvic kidney: A case report. Int J Surg Case Rep 2020; 66:115-117. PMID: 31837612 DOI: 10.1016/j.ijscr.2019.11.041.
17. Robertson AG, Kim J, Al-Ahmadie H, Vitaly Margulis , Jay Raman , Yair Lotan , Timothy O'Brien , Nancy You , Shahrokh F Shariat, Surena F Matin Comprehensive molecular characterization of muscle-invasive bladder cancer. Cell 2017; 171(3):540–56.e25. PMID: 25711197 DOI: 10.1016/j.juro.2015.02.081.
18. Roupert M, Babjuk M, Compérat E, Zigeuner R, Sylvester RJ, Burger M, Nigel C, Cowan NC, Böhle A, Rhijn BWG, Kaasinen K, Palou J, Shariat SF . European Association of Urology guidelines on upper urinary tract urothelial cell carcinoma: 2015 update. Eur Urol 2015; 68: 868–879. PMID: 26188393 DOI: 10.1016/j.eururo.2015.06.044.
19. Roupert M, Yates DR, Comperat E, Cussenot O. Upper urinary tract urothelial cell carcinomas and other urological malignancies involved in the hereditary nonpolyposis colorectal cancer (lynch syndrome) tumor spectrum. Eur Urol 2008; 54(6):1226–1236. PMID: 18715695 DOI: 10.1016/j.eururo.2008.08.008.
20. Sijmons RH, Kiemeneij LA, Witjes JA, Vasen HF. Urinary tract cancer and hereditary nonpolyposis colorectal cancer:

- risks and screening options. J Urol 1998; 160: 466–470. PMID: 9679899.
21. Van der Post RS, Kiemeny LA, Ligtenberg MJ, Hulsbergen-van de Kaa CA, Bodmer L, Schaap CM, Kets JHJM van Krieken, Hoogerbrugge N. Risk of urothelial bladder cancer in Lynch syndrome is increased, in particular among MSH2 mutation carriers. J Med Genet 2010; 47:464–47. DOI: 10.1136/jmg.2010.076992.
  22. Vikas S, Mona B, Smita G, Rahul Y, Shazib F, Payel D, Dinesh S, Ankur A, Rafey K, Kshitij M. Urothelial carcinoma in young adults-Single centre experience. Medical Science, 2020, 24(106), 3967-3971